## Prime Farmland

(Only the soils considered prime farmland are listed. Urban or built-up areas of the soils listed are not considered prime farmland. If a soil is prime farmland only under certain conditions, the conditions are specified in parentheses after the soil name.)

## Worth County, Iowa

	T
Map	Soil name
symbol	 
27	  Terril loam, 2 to 5 percent slopes
29	Clarion-nicollet loams, 1 to 3 percent slopes
55	Nicollet loam, 1 to 3 percent slopes
83	Kenyon loam, 0 to 2 percent slopes
83B	Kenyon loam, 2 to 5 percent slopes
84	Clyde silty clay loam, 0 to 2 percent slopes (Prime farmland if drained)
95	Harps loam, 1 to 3 percent slopes (Prime farmland if drained)
107 135	Webster silty clay loam, 0 to 2 percent slopes (Prime farmland if drained)  Coland silty clay loam, 0 to 2 percent slopes (Prime farmland if drained and either protected   from flooding or not frequently flooded during the growing season)
138	Clarion loam, 0 to 2 percent slopes
138B	Clarion loam, 2 to 5 percent slopes
151	Marshan clay loam, moderately deep, 0 to 2 percent slopes (Prime farmland if drained)
152	Marshan clay loam, deep, 0 to 2 percent slopes (Prime farmland if drained)
153	Marshan clay loam, depressional, 0 to 1 percent slopes (Prime farmland if drained)
169B	Clarion loam, 2 to 5 percent long slopes
171	Bassett loam, 0 to 2 percent slopes
171B	Bassett loam, 2 to 5 percent slopes
173	Hoopeston fine sandy loam, 0 to 2 percent slopes
174	Bolan loam, 0 to 2 percent slopes
174B	Bolan loam, 2 to 5 percent slopes
175 175B	Dickinson fine sandy loam, 0 to 2 percent slopes  Dickinson fine sandy loam, 2 to 5 percent slopes
1775	Saude loam, 0 to 2 percent slopes
177B	Saude loam, 2 to 5 percent slopes
178	Waukee loam, 0 to 2 percent slopes
178B	Waukee loam, 2 to 5 percent slopes
184	Klinger silty clay loam, 1 to 3 percent slopes
188	Kensett silt loam, 0 to 2 percent slopes
198	Floyd loam, 1 to 3 percent slopes
213	Rockton loam, deep, 0 to 2 percent slopes
213B	Rockton loam, deep, 2 to 5 percent slopes
214	Rockton loam, moderately deep, 0 to 2 percent slopes
214B	Rockton loam, moderately deep, 2 to 5 percent slopes
225	Lawler loam, moderately deep, 0 to 2 percent slopes
226 235	Lawler loam, deep, 0 to 2 percent slopes  Coland-turlin complex, 0 to 2 percent slopes (Prime farmland if drained and either protected
233	from flooding or not frequently flooded during the growing season)
236B	Lester loam, 2 to 5 percent slopes
236B2	Lester loam, 2 to 5 percent slopes, moderately eroded
325	Le sueur loam, 1 to 3 percent slopes
329	Webster-nicollet complex, 1 to 3 percent slopes (Prime farmland if drained)
335	Harcot loam, 0 to 2 percent slopes (Prime farmland if drained)
377	Dinsdale silty clay loam, 0 to 2 percent slopes
377B	Dinsdale silty clay loam, 2 to 5 percent slopes
382	Maxfield silty clay loam, 0 to 2 percent slopes (Prime farmland if drained)
399	Readlyn loam, 1 to 3 percent slopes
407	Schley silt loam, 1 to 3 percent slopes (Prime farmland if drained)
471	Oran silt loam, 1 to 3 percent slopes (Prime farmland if drained)
507	Canisteo silty clay loam, 0 to 2 percent slopes (Prime farmland if drained)
558 559	Talcot clay loam, moderately deep, 0 to 2 percent slopes (Prime farmland if drained)  Talcot clay loam, deep, 0 to 2 percent slopes (Prime farmland if drained)
583	Minnetonka silty clay loam, 1 to 3 percent slopes (Prime farmland if drained)
638B2	Clarion-storden loams, 2 to 5 percent slopes, moderately eroded
	· · · · · · · · · · · · · · · · · · ·

USDA-NRCS, IOWA 04/10/2002

## Prime Farmland

(Only the soils considered prime farmland are listed. Urban or built-up areas of the soils listed are not considered prime farmland. If a soil is prime farmland only under certain conditions, the conditions are specified in parentheses after the soil name.)

## Worth County, Iowa

Map symbol	Soil name
695	Tilfer silty clay loam, 0 to 2 percent slopes (Prime farmland if drained and either protected
706	from flooding or not frequently flooded during the growing season)
706 706B	Donnan silt loam, dark variant, 0 to 2 percent slopes  Donnan silt loam, dark variant, 2 to 5 percent slopes
706B 725	Hayfield loam, moderately deep, 0 to 2 percent slopes
726	Hayfield loam, deep, 0 to 2 percent slopes
733	Calco silty clay loam, loamy substratum, 0 to 2 percent slopes (Prime farmland if drained and   either protected from flooding or not frequently flooded during the growing season)
755	Nicollet loam, 1 to 3 percent long slopes
761	Franklin silt loam, 1 to 3 percent slopes
777	Wapsie loam, 0 to 2 percent slopes
777B	Wapsie loam, 2 to 5 percent slopes
782	Donnan silt loam, 0 to 2 percent slopes
782B	Donnan silt loam, 2 to 5 percent slopes
836B	Kilkenny silty clay loam, 2 to 5 percent slopes
836B2	Kilkenny silty clay loam, 2 to 5 percent slopes, moderately eroded
855	Shorewood silty clay loam, 1 to 3 percent slopes
977	Richwood silt loam, 0 to 2 percent slopes

USDA-NRCS, IOWA 04/10/2002